

Brief Reports

The Effect of Audiovisual Distraction on Pain Perception During Reduction of Distal Radial Fractures: A Dual Center Randomized Controlled Trial

Hilmar H. Heijmans, MD,^{*,#} Maybritt I. Kuypers, MD, PHD,^{*,#} Milan L. Ridderikhof, MD, PHD,^{*} Kaoutar Azijli, MD, PHD,[†] Susan van Dieren, PHD,^{‡,§} and Marlies P. Schijven, MD, PHD^{‡,§,||}

^{*}Department of Emergency Medicine, Amsterdam UMC, Location AMC, Amsterdam, The Netherlands, [†]Department of Emergency Medicine, Amsterdam UMC, Location VUMC, Amsterdam, The Netherlands, [‡]Department of Surgery Amsterdam UMC, University of Amsterdam, Amsterdam, The Netherlands, [§]Amsterdam Gastroenterology and Metabolism AGEM, Amsterdam, The Netherlands, and ^{||}Amsterdam Public Health program Digital Health APH, Digital Health, Amsterdam, The Netherlands

Reprint Address: H.H. Heijmans, MD, Department of Emergency Medicine, Amsterdam UMC location AMC, Meibergdreef 9, Amsterdam, The Netherlands

Abstract—Background: Adequate pain management is a challenge in the Emergency Department (ED). This study explores whether audiovisual distraction (AVD) has a positive effect on pain perception. **Objective:** We aimed to compare the influence of AVD on reported pain, anxiety and satisfaction using a commercial AVD device as an adjunct to standard care in patients undergoing distal radial fracture reduction in the ED. **Methods:** This randomized controlled, dual-center study, conducted in the ED of 2 Level-1 trauma centers, included a total of 36 adult patients with a distal radial fracture requiring reduction in the ED. Patients were randomly assigned to either standard of care or the use of AVD in addition to standard of care. **The primary outcome was reported pain after fracture reduction using a standardized verbal Numerical Rating Scale (v-NRS). Results:** No significant differences were found between groups in pain (difference: 0.8; 95% CI, -1.2 to 2.8), anxiety (difference: 0.6; 95% CI, -1.6 to 2.8) or satisfaction (difference: 0.1; 95% CI, -0.7 to 1.0). However, the majority (>80%) of the patients, nurses and physicians stated they would recommend the use of AVD for future procedure. **Conclusion:** This study was unable to demonstrate the added value of using audiovisual immersion as a distraction tool to standard care, in terms of lowering perceived pain or anxiety in adults undergoing a wrist fracture reduction in the ED.

Nevertheless, patients and healthcare providers appeared to be satisfied and would recommend the use of the AVD device. © 2025 The Author(s). Published by Elsevier Inc. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>)

Keywords—audiovisual distraction; pain; anxiety; fracture; virtual reality

Introduction

Adequate and safe relief of acute pain during painful procedures remains a major global challenge in Emergency Care (1). More than half of all patients presenting to the Emergency Department (ED) report experiencing pain (2). To date, peri-procedural pain management for the treatment of acute pain in the ED setting remains sub-optimal (3).

One of the oldest non-pharmacological methods known to be effective and safe to mitigate pain is mental distraction (4,5). With the rise of high-tech video glasses and virtual reality (VR), distraction techniques have re-gained increasing attention as an adjunct for use in healthcare. Specifically, research has shown VR to be promising in the field of pain management. This includes

[#]shared first authorship.

wound care of burns in children, cystoscopy, and various surgical and dental procedures (6–9). Moreover, audiovisual distraction (AVD) has been shown to have potential to decrease sedative dosages needed in pain management (10).

How AVD works to reduce pain is only partially known. The gate control theory postulates pain perception to be modulated by interaction among different neurons. AVD or other stimuli might lead to the closure of certain neural gateways, thus reducing pain perception (11). Virtual reality (VR) appears to be more effective than traditional methods due to its immersive properties (12). It demands engagement from a patient's visual, auditory and, in some cases, also physical actions.

Distal radius fractures are the most common upper extremity fractures presented in the ED (13). When displaced, they can be repositioned manually or with the help of finger-trap traction. Reduction of the fracture is known to be a painful procedure, frequently performed in the ED (14). Placing an inter-osseous hematoma block in the fracture gap using lidocaine is a frequently used technique to obtain analgesia for this type of fracture reduction and is considered to be a safe and effective method (15). The analgesic effect of the hematoma block depends on the technique of injection and the amount of lidocaine used (16).

Our aim was to investigate whether the additional application of a commercial portable multimedia device with audiovisual glasses (HappyMed®) would be effective in reducing pain in adults with a displaced wrist fracture in need for manual reduction in the ED. We hypothesized that the addition of this AVD device to the standard of care (SOC) would result in lower pain scores, less anxiety and higher patient satisfaction. Additionally, our objective was to explore the feasibility of implementing a new innovative tool in a busy ED by evaluating the satisfaction and feedback from the emergency physicians and nurses.

Methods

Study Design

A randomized controlled trial was conducted in 2 Dutch academic level-1 trauma centers to compare SOC with SOC augmented with AVD. The annual ED census of Amsterdam UMC location AMC was 22483 and Amsterdam UMC location VUMC was 20713. The study was conducted in accordance with the CONSORT guidelines (17). The protocol was approved by the Institutional Review Board of the relevant study sites (Medisch Ethische Toetsing Commissie – Amsterdam UMC location AMC - NL67837.018.18). All included patients provided written informed consent.

Participants

Eligible patients were adults presenting to the ED with a displaced distal radial fracture requiring manual reduction, after assessment of radiographic imaging, and who presented within 24 hours after the sustained injury. Exclusion criteria were visual or auditory impairment, multiple traumatic injuries, illiterate, non-Dutch speaking, a known history of anxiety disorder, intoxication, chronic opioid use, lidocaine allergy, or an indication for acute surgical intervention. Study participants were patients who met the eligibility criteria and were both willing and able to give informed consent.

Recruitment

Patients underwent initial assessment and management by an ED triage nurse. Pain management was implemented according to the standard hospital protocols. Board-certified emergency physicians, who would also perform the hematoma block and reduction, approached the patients and provided verbal and written information about the study. If the patient was interested, additional questions were answered. Patients were fully assessed against the inclusion and exclusion criteria before written consent was obtained for those willing and able to participate.

Study Procedure and Data Collection

Participants in both groups (SOC and SOC + AVD) received instructions on how to grade their pain, level of anxiety and satisfaction using the v-NRS; an 11-point scale ranging from 0 to 10 which is validated for measuring acute pain in the ED (18). With 0 indicating “no pain,” “no anxiety” or “not satisfied at all with the total procedure” and 10 indicating “the worst pain possible,” “severe anxiety” or “completely satisfied with the total procedure.”

Pain perception and anxiety were measured right before the start of the procedure (before the administration of the hematoma block) and immediately after (when the plaster cast had been applied). When in cast (e.g., after reduction), patients were asked to score the maximum level of pain and anxiety they had experienced during the administration of the hematoma block and the fracture reduction. Just before discharge, patients were asked to score their overall satisfaction with the procedure. The patients in the intervention group (SOC+AVD) were asked to rate the quality of the audio and visual effects and the distraction on an 11-point scale and whether they would use the AVD device again under similar circumstances.

Data were collected on a standardized case report form by the treating physician and included various patient

characteristics and questions; these can be found in supplement A and B.

The informed consent and the case report forms were stored in an opaque envelope and the data was entered in a dedicated, password protected, SSL-encrypted database (Castor®, Ciwit BV, Amsterdam, The Netherlands) by the investigators.

Interventions

The standard operation procedure was taught to all Emergency Physicians involved at the study sites. This was included in a package with study information in addition to consent forms in a concealed study envelope. Participants in both groups received similar care with regard to analgesic block and reduction method. All participants received the same information on the procedural steps (first, hematoma block; second, 10 min finger traction; third, manual reduction; and lastly, a plaster cast). A hematoma block was performed by injecting lidocaine into the fracture hematoma. A standard dosage of 20 mL of lidocaine 1% for adult patients with a body weight >50 kilograms was advised. Specific dosages of lidocaine were prescribed for patients weighing <50 kilograms. For increased distribution and analgesic effect, it was advised to inject 10 mL into the hematoma directly after successful withdrawal of blood in the syringe. Subsequently 10 mL was divided between the lateral and medial surface of the fractured radius. Hereafter, a finger trap traction device was applied and used for providing continuous fracture traction for a period of 10 minutes. Patients were asked to report their pain to ascertain the analgesic effect of the hematoma block before reducing the fracture. In the case that patients reported a pain score (v-NRS) >6, additional analgesia was offered. Consequently, the fracture was reduced manually, and a plaster cast was applied. Finally, a radiographic image in 2 dimensions was performed to check the fracture angles after reduction.

In addition to the SOC, a portable multimedia device was used in patients assigned to the intervention group. This device, referred to as the 'HappyMed®', consists of a pair of video glasses with headphones which is connected to a controller with a touchscreen. The headset is adjustable and shields the patient off completely from their surroundings. The high-quality video and audio are intended to give the patient a 'real' cinematic experience. In contrast to VR, HappyMed® video glasses show only 2-dimensional content. This is to ensure patients do not move their head/body as this may cause unintentional additional pain or problems with the procedure.

Prior to administration of the hematoma block, the AVD device was given to the patient and a movie of choice was started. [Figure 1](#) shows a picture of the set-up in the intervention group. Five minutes after the initiation of the




Figure 1. Demonstration of the audiovisual glasses.

movie, the hematoma block was given. The AVD device was first removed after immobilizing the wrist with a plaster cast.

Randomization, Allocation Concealment, and Blinding

Randomization with a 1:1 ratio to either intervention (SOC+AVD) or control group (SOC) was done via a secure web-based randomization system (Castor®). The treating emergency physician accessing the randomization website did not know the allocation for an individual patient until recruitment was confirmed. Blinding was not possible for this study due to the nature of the intervention.

Primary Outcome

The primary outcome of this study was the maximum pain experienced during placement of the hematoma block and the manual fracture reduction.

Secondary Outcomes

Our secondary outcomes included patient anxiety and satisfaction scores, adverse events, analgesic medication used, success rate of the manual fracture reduction and

satisfaction of nurses and emergency physicians on the use of the AVD device.

Sample Size

The main objective of this study was to assess if there is a clinically significant difference in maximum pain intensity scores during a painful procedure with and without audiovisual immersive technology. Using the criteria of a balanced sensitivity and specificity, the best cut-off points determined by Farrar et al. (19) for pain intensity are $>$ or $=2$ difference on a 0-10 numeric rating scale. Assuming a standard deviation of 2, and using a 2 tailed 2 sample t test, with a type 1 error rate of 0.05, a sample size of 17 participants per group provides sufficient power (80%) to detect a between group difference of 2 or more points on the v-NRS.

Statistical Analyses

The data were analyzed using an intention-to-treat analysis. Baseline characteristics were presented using descriptive statistics. Continuous data were expressed as means with standard deviations and categorical data as numbers and percentages. Normality of the data distribution was analyzed by visually inspecting the histograms. For the primary and secondary outcomes, according to data distribution, a 2-sided Student's t-test or Mann Whitney U test was used. The mean difference between the 2 groups was presented together with 95% confidence intervals (CI). The secondary outcomes of anxiety and patient satisfaction scores were also examined for normality and tested using either the students t-test or the Mann-Whitney U test. The differences between the groups were displayed as mean difference with 95% CI or interquartile range (IQR).

Results

Recruitment and Flow Through Trial

Figure 2 outlines the flow of participants in our study. Recruitment took place from June 2019 to February 2021. Ten patients were excluded: 5 did not speak Dutch, 3 patients suffered from a multi-trauma and 2 patients opted out as they preferred to be able to see the procedure. Thirty-seven patients entered the study after providing informed consent and were randomly assigned: 18 to the control and 19 to the intervention group. Two patients could not participate: 1 because of a malfunctioning AVD device and another due to an error of the randomization software. When the AVD device malfunctioned for a second time, we changed this patient from intervention to

control group. Hereby leaving 17 patients in the intervention group and 19 in the control group.

Baseline Characteristics

Baseline characteristics are shown in Table 1. The patients ranged in age from 21 to 86 years with a mean of 56 years. There were no significant differences in groups for age, sex, fracture type, baseline pain and baseline anxiety. However, the control group reported a higher baseline pain (6.2 vs 5.5 $p = 0.36$) and lower anxiety score (3.7 vs 4.9 $p = 0.20$) compared with the intervention group. This was not considered statistically significant. Administration of pre-hospital pain medication was comparable in both groups. Although the recommended dosage of lidocaine was prescribed in the standard operating procedure, the doses administered varied from 8 mL to 20 mL. No significant difference was found in dosage between the groups. Fentanyl was given intravenously in 2 patients in the control group and in 1 patient in the intervention group.

Primary Outcome

No significant differences in pain during analgesic injection or fracture reduction were found between the control group and intervention group (Table 2). The mean reported pain during injection was higher in the control group: 5.3 (SD 2.4) vs 4.4 (SD 2.5), $p = 0.31$. The mean reported pain for the fracture reduction was higher in the intervention group (4.8; SD 3.0 vs 5.6; SD 3.0 $p = 0.43$) (Figure 3).

Secondary Outcomes

No significant difference between the groups was found in reported anxiety or satisfaction scores. (Table 2 and Figure 4). Patients in the intervention group graded the quality of the video with an average of 7.8 and the audio with a 7.9 on a 10-point scale (1-10). The effect of the distraction of the video glasses was given a 6.2 out of 10. When asked if they would use the video glasses next time, 83% of the patients, 88% of the physicians and 94% of the nurses answered with a 'yes'. One physician reported that the AVD device negatively influenced the procedure, without further specification, and another commented that the intervention "took too much time". One patient reported that "the volume of the audio was too low". There were no adverse events reported.

Discussion

The current randomized controlled trial (RCT) suggests that the application of an AVD device, in ad-




Figure 2. CONSORT flow diagram describing the patient selection process.
SOC: standard of care; CONSORT: Consolidated Standards of Reporting Trials

Table 1. Baseline Characteristics.

Characteristics	Control group (n = 19)	Intervention group (n = 17)
Age in years, mean (SD)	57.5 (18.0)	54.7 (20.2)
Sex, N (%)		
Male	3 (15.8)	5 (29.4)
Female	16 (84.2)	12 (70.6)
Fracture type, N (%)		
Dorsal	15 (78.9)	12 (70.6)
Volar	4 (21.1)	5 (29.4)
Baseline Pain-Score (SD)	6.2 (1.6)	5.5 (2.9)
Baseline Anxiety-Score (SD)	3.7 (2.6)	4.9 (2.7)
Pre-hospital medication, N (%)		
Acetaminophen 1000 mg	10 (52.6)	10 (58.8)
Diclofenac 50 mg	5 (26.3)	2 (11.8)
Ibuprofen 400 mg	0	1 (5.9)
Pre-procedural medication, N (%)		
Acetaminophen 1000 mg	6 (31.6)	1 (5.9)
Diclofenac 50 mg	3 (15.8)	1 (5.9)
Fentanyl 1 µg/kg	2 (10.5)	1 (5.9)
Piritramide 15 mg	0	1 (5.9)
Procedural medication		
Lidocaine 1% in mL, mean (SD) Additional medication during procedure, N (%)	13.2 (4.7)	10.3 (4.4)
Fentanyl 1 µg/kg	1 (5.3)	1 (5.9)
Piritramide 15 mg	1 (5.3)	0

SD: standard deviation.




Figure 3. Mean v-NRS pain scores during analgesic injection and during fracture reduction. v-NRS: verbal numeric rating scale.

Table 2. Primary and Secondary Outcomes.

Variable	Control group (n = 19)	Intervention group (n = 17)	Difference (95% CI)	P
Maximum pain score (v-NRS) during:				
Analgesic injection	5.3 (2.4)	4.4 (2.5)	-0.9 (-2.5 to 0.8)	0.31
Fracture reduction	4.8 (3.0)	5.6 (3.0)	0.8 (-1.2 to 2.8)	0.43
Anxiety score	3.5 (3.0)	4.1 (3.4)	0.6 (-1.6 to 2.8)	0.59
Satisfaction score				
Patient	8.6 (1.3)	8.7 (1.3)	0.1 (-0.7 to 1.0)	0.77
Physician	8.1 (0.7)	7.4 (1.4)	-0.7 (-1.5 to 0.1)	0.07
Nurse	8.2 (1.0)	8.0 (1.4)	-0.2 (-1.1 to 0.7)	0.65
Would recommend HappyMed®				
Patient	-	83% (14/17)		
Physician	-	88% (14/16)		
Nurse	-	94% (16/17)		
Score HappyMed®				
Quality of video	-	7.8		
Quality of audio	-	7.9		
Distraction	-	6.2		
Fracture reduction successful after first attempt	73% (11/15)	86% (12/14)		
Adverse events	0	0		

CI = confidence interval; v-NRS = verbal numerical rating scale; SD = standard deviation.

dition to the SOC does not provide sufficient distraction to significantly reduce pain and anxiety in patients undergoing reduction of a distal radial fracture in the ED. These findings contradict earlier reported studies in literature, in which the authors showed a pain and anxiety reducing effect when using audiovisual immersion using VR to complement standard care (20,21).

One of the reasons may be that these studies used VR, which has more immersive qualities compared with the HappyMed® AVD device. According to Sinha et al., the greater the immersion the greater the analgesic effect will be. (22) The score for overall distraction in our study was rated with an average of 6. This sits in contrast to the 8+ score which was given for the quality of the video and audio of HappyMed®. It might be that the HappyMed® headset is satisfactory for watching a movie but is not sufficient in providing enough distraction during painful procedures. We know from other studies that have used VR, that the time spent thinking about pain during medical procedures is significantly less compared to watching television as distraction. (23,24) However, a recent investigation published in 2023, compared 2-dimensional (2D)

audio-video experiences with 3-dimensional (3D) virtual reality in adult emergency department patients requiring a minor procedure. (25) Distraction by viewing a 3D virtual world in a head-mounted VR display did not result in lower average levels of procedural pain and anxiety than that by 2D viewing on a screen, despite a higher sense of telepresence. Another important reason for the differing results may be that the aforementioned VR studies were conducted in predominantly pediatric or young adolescent patients. Again, this sits in contrast to our predominantly adult female population with a mean age of 56. Children are well known to be more susceptible to immersive and distractive techniques when compared to adults. (26) For example, Olesen *et al.* demonstrated that children exhibit distinct brain activation patterns in response to distraction tasks, suggesting they are inherently more susceptible to such interventions. (25) Similarly, Birnie *et al.* provided evidence from a systematic review that distraction significantly reduces procedural pain and distress in pediatric patients, of finding that aligns earlier work by McCaul and Malott on the overall effectiveness of distraction in modulating pain perception. (5,26) However, these benefits observed in children may not directly




Figure 4. Mean anxiety and satisfaction v-NRS scores in patients receiving SOC with or without audiovisual distraction using the HappyMed device. v-NRS: verbal numeric rating scale; SOC: standard of care.

translate to older adults, given the differences in cognitive processing, attention and pain perception between age groups. Although not statistically significant, we found that physicians rated satisfaction higher in the control group. This was also reflected in some of the reported comments provided by physicians. Some noted the additional intervention was too time-consuming. This is an important point. The ED can be a busy place where time and personnel are scarce at times. There may be threshold for implementing new devices if they appear to take too much time to set up. According Birrenbach *et al.* (27) VR seems to be effective and implementable in the ED. They enrolled fifty-two patients mostly with pain in extremities ($n = 15$, 28.8%) and abdomen ($n = 12$, 23.1%). It's important to note that this was a convenience sample. As such, it is unclear if VR was, for example, also implemented during a busy ED shift. About 1 third of patients in this study presented with trauma-associated pain ($n = 16$, 30.8%). When compared to our study, all patients presented with traumatic pain, requiring a painful procedure. The duration of pain in their study was subacute (> 24 h) ($n = 32$, 61.5%) in most patients. In our study all patients presented with acute pain (< 24 hrs). These differences may be other indicators as to why we found different results.

In the study of Birrenbach *et al.* the authors concluded they found a significant reduction in pain (NRS median pre-VR simulation 4.5 (IQR 3–7) vs. median post-VR simulation 3 (IQR 2–5), $p < 0.001$). It is debatable whether a reduction from 4.5 to 3 (a difference of 1.5) is a clinically relevant reduction in pain. In our study we considered ≥ 2 -point difference on the NRS scale to be clinically significant; this was based on a validation study examining the changes in acute pain mea-

surement scales that are most strongly associated with a patient-determined indicator of clinical importance. (19) Clinically significant pain reduction is an important factor to consider for future studies.

Despite the lack of a statistically significant reduction in pain or anxiety, patients appeared to derive enjoyment and comfort from watching a movie during the procedure. For instance, 1 patient specifically requested to continue watching for a longer period, and another wished to wear the video glasses again for a second reduction attempt, even though it was not part of the study. These experiences suggest that the device offered a positive overall patient-experience, which could explain why patients, nurses and physicians all recommended the use of the intervention.

Limitations

There were some differences between the 2 groups, including baseline pain and anxiety scores as well as dosing of analgesic and anesthetic medications. Although these differences were not statistically significant, we cannot rule out their influence on the study outcomes. Additionally, due to the limited sample size, it is possible that procedural complexity varied between the groups. Nuances in the technique of performing the hematoma block or the reduction procedure, may have led to unmeasured differences.

We encountered issues with the recruitment of subjects. Although wrist fracture reduction is a common presentation in the ED, the presentations were very low in the initial research location. Our study sites are level 1 trauma centers and mostly receive multi-trauma patients. Wrist fractures are more frequently seen in level 2 or 3 trauma centers in the Netherlands. Fur-

ther delay was caused by the outbreak of the COVID-19 pandemic, in this timeframe all clinical research was temporarily ceased; this caused significant delay in recruitment.

Our study design aimed to replicate a real-world clinical scenario where the audiovisual distraction is fully implemented. Participants were free to choose from over 100 movie titles on the HappyMed™ device. We believe that they are likely to be more engaged and distracted by a movie they find appealing, compared to one that does not capture their interest. While this patient-centered approach enhances the experience, it may introduce variability in the level of distraction. Blinding the physician was not possible and thus could lead to observer bias. Although blinding using an ‘off’ headset for the control group was considered, we were concerned that it might alter patient behavior and inadvertently increase their focus on pain. To mitigate selection bias, randomization and allocation were concealed through a computer program after assessment of eligibility and inclusion.

Conclusion

This study did not find a clinically significant difference in pain or anxiety in patients using an AVD device during distal radial fracture reduction in the ED. While objective measures of pain and anxiety did not differ significantly between groups, satisfaction was equally high and willingness to use the device in future procedures in the intervention group was also high. Due to our study limitations, we cannot draw any definite conclusions. Further investigations are warranted to explore the potential benefits and negative side-effects of AVD, especially in pediatric and elderly patients, in the emergency department setting. We recommend future research to focus on clinically relevant differences in pain. Lastly, it is important to consider that an innovative tool needs to be intuitive and easy to implement and maintain in the ED setting.

Funding

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Supplementary materials

Supplementary material associated with this article can be found, in the online version, at [doi:10.1016/j.jemermed.2025.03.015](https://doi.org/10.1016/j.jemermed.2025.03.015).

CRedit authorship contribution statement

Hilmar H. Heijmans: Writing – review & editing, Writing – original draft, Project administration, Investigation, Formal analysis, Data curation. **Maybritt I. Kuypers:** Writing – review & editing, Writing – original draft, Supervision, Methodology, Conceptualization. **Milan L. Ridderikhof:** Writing – review & editing, Supervision. **Kaoutar Azijli:** Investigation. **Susan van Dieren:** Methodology, Formal analysis. **Marlies P. Schijven:** Supervision, Software, Conceptualization.

References

1. Upp J, Kent M, Tighe PJ. The evolution and practice of acute pain medicine. *Pain Med. (United States)* 2013;14:124–44.
2. Berben SAA, Meijs THJM, van Dongen RTM, et al. [Guideline ‘Pain management for trauma patients in the chain of emergency care’]. *Ned. Tijdschr. Geneesk.* 2011;155:A3100.
3. Arendts G, Fry M. Factors associated with delay to opiate analgesia in emergency departments. *J. Pain* 2006;7:682–6.
4. Blount RL, Zempsky WT, Jaaniste T, et al. *The Guilford Press*; 2009. p. 171–88.
5. McCaul KD, Malott JM. Distraction and coping with pain. *Pain* 1984;23:315.
6. Hoffman HG, Chambers GT, Meyer WJ, et al. Virtual reality as an adjunctive non-pharmacologic analgesic for acute burn pain during medical procedures. *Ann. Behav. Med.* 2011;41:183–91.
7. Jeffs D, Dorman D, Brown S, et al. Effect of virtual reality on adolescent pain during burn wound care. *J. Burn Care Res.* 2014;35:395–408.
8. Mosso JL, Gorini A, De La Cerda G, et al. Virtual reality on mobile phones to reduce anxiety in outpatient surgery. *Studies in Health Technology and Informatics* 2009;142.
9. Walker MR, Kallingal GJS, Musser JE, et al. Treatment efficacy of virtual reality distraction in the reduction of pain and anxiety during cystoscopy. *Mil. Med.* 2014;179:891–6.
10. Lee DWH, Chan ACW, Wong SKH, et al. Can visual distraction decrease the dose of patient-controlled sedation required during colonoscopy? A prospective randomized controlled trial. *Endoscopy* 2004;36:197–201.
11. Van Griensven H. *Pain: a textbook for health professionals*; 2014.
12. Greesma SG, George S, Anandaraj S, et al. Comparative evaluation of the efficacy of virtual reality distraction, audio distraction and tell-show-do techniques in reducing the anxiety level of pediatric dental patients: an In vivo study. *Int. J. Clin. Pediatr. Dent.* 2021;14:S173–8.
13. Shauver MJ, Clapham PJ, Chung KC. An economic analysis of outcomes and complications of treating distal radius fractures in the elderly. *J. Hand Surg. Am.* 2011;36 1912–1918.e3.
14. Padegimas EM, Ilyas AM. Distal radius fractures. *Emergency Department Evaluation and management. Orthop. Clin. North Am.* 2015;46:259–70.

15. Tseng PT, Leu TH, Chen YW, Chen YP. Hematoma block or procedural sedation and analgesia, which is the most effective method of anesthesia in reduction of displaced distal radius fracture? *J. Orthop. Surg. Res.* 2018;13:1–7.
16. Orbach H, Rozen N, Rinat B, Rubin G. Hematoma block for distal radius fractures – Prospective, randomized comparison of two different volumes of lidocaine. *J. Int. Med. Res.* 2018;46:4535–8.
17. Barbour V, Bhui K, Chescheir N, et al. CONSORT Statement for randomized trials of nonpharmacologic treatments: A 2017 update and a CONSORT extension for nonpharmacologic trial abstracts. *Ann. Intern. Med.* 2017;167:40–7.
18. Bijur PE, Latimer CT, Gallagher EJ. Validation of a verbally administered numerical rating scale of acute pain for use in the emergency department. *Acad. Emerg. Med.* 2003;10:390–2.
19. Farrar JT, Berlin JA, Strom BL. Clinically important changes in acute pain outcome measures: A validation study. *J. Pain Symptom Manage.* 2003;25:406–11.
20. Chan E, Foster S, Sambell R, Leong P. Clinical efficacy of virtual reality for acute procedural pain management: A systematic review and meta-analysis. *PLoS One* 2018;13:1–13.
21. Zhang C, Qin D, Shen L, Ji P, Wang J. Does audiovisual distraction reduce dental anxiety in children under local anesthesia? A systematic review and meta-analysis. *Oral Dis* 2019;25:416–24.
22. Sinha M, Christopher NC, Fenn R, Reeves L. Evaluation of non-pharmacologic methods of pain and anxiety management for laceration repair in the pediatric emergency department. *Pediatrics* 2006;117:1162–8.
23. Hoffman HG, Rodriguez RA, Gonzalez M, et al. Immersive virtual reality as an adjunctive non-opioid analgesic for pre-dominantly Latin American children with large severe burn wounds during burn wound cleaning in the Intensive care unit: a pilot study. *Front. Hum. Neurosci.* 2019;13.
24. Spiegel B, Fuller G, Lopez M, et al. Virtual reality for management of pain in hospitalized patients: a randomized comparative effectiveness trial. *PLoS One* 2019;14:1–15.
25. Olesen PJ, Macoveanu J, Tegnér J, Klingberg T. Brain activity related to working memory and distraction in children and adults. *Cereb. Cortex* 2007;17:1047–54.
26. Birnie KA, Noel M, Parker JA, et al. Systematic review and meta-analysis of distraction and hypnosis for needle-related pain and distress in children and adolescents. *J. Pediatr. Psychol.* 2014;39:783–808.
27. Birrenbach T, Bühlmann F, Exadaktylos AK, et al. Virtual Reality for Pain Relief in the Emergency Room (VIPER) - a prospective interventional feasibility study. *BMC Emerg Med.* 2022;22:113.

Article Summary

1. Why is this topic important?

Pain management in the Emergency Department remains suboptimal. This study explores a new innovative method to reduce pain during medical procedures.

2. What does this study attempt to show?

This study attempts to show that audiovisual distraction with the help of video glasses for patients with a wrist fracture can be an effective and feasible non-pharmacological pain treatment.

3. What are the key findings?

No statistically significant differences were found in pain, anxiety and satisfaction scores between both groups. However the method seems safe and feasible.

4. How is patient care impacted?

We tested audiovisual distraction during wrist fracture reduction to improve patient experience. However this study has not found a difference in patient reported outcomes. Thus, patient care has not (yet) been impacted.